CHLORAMIN

Water Soluble Powder

Composition:

Each sachet of 100 g contains:

Chlortetracycline hydrochloride 20 g

Description:

Chlortetracycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria like Bordetella, Campylobacter, Chlamydia, E. coli, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. The action of chlortetracycline is based on inhibition of bacterial protein synthesis. Chlortetracycline is mainly excreted in urine, for a small part in bile and in lactating animals in milk.

Indications:

Gastrointestinal and respiratory infections caused by Chlortetracycline sensitive bacteria in cattle, sheep, goats and poultry .

Contraindications:

- Hypersensitivity to tetracyclines.
- Animals with renal or hepatic failure .

Side effects:

Hypersensitivity reaction.

Discoloration of teeth in young animals.

Usage:

Oral with drinking water

Dosage:

Cattle: 40 g per animal for 3-5 days

Sheep, Goats: 10 g per animals for 3 days.

Poultry: 100-200 g per 200 liter of drinking water for 3-5 days.

Withdrawal Period:

Ruminants: Meat: 8 days Milk: 3 days.

Poultry: Meat :3 days eggs : 5 days

Storage conditions:

Store in airtight containers, in a moderate temperature (15-25 C) and protect

form direct sunlight.

Packing:

Sachet of 100 g.